

Tamaño del Pene

Porcentaje de hombres que tienen el pene erecto más pequeño que el tamaño indicado

Medida (cm)	Medida (pulgadas)	Estudio Koreano Longitud	Estudio Mexicano Longitud	Estudio Mexicano Circunferencia
5.0	2.0	0.0%	0.0%	0.6%
5.5	2.2	0.0%	0.0%	1.1%
6.0	2.4	0.1%	0.0%	1.7%
6.5	2.6	0.1%	0.0%	2.2%
7.0	2.8	0.2%	0.0%	2.8%
7.5	3.0	0.5%	0.0%	3.3%
8.0	3.1	0.8%	0.1%	3.7%
8.5	3.3	1.5%	0.1%	4.6%
9.0	3.5	2.5%	0.2%	5.7%
9.5	3.7	4.1%	0.5%	8.1%
10.0	3.9	6.4%	1.0%	10.4%
10.5	4.1	9.6%	1.8%	16.7%
11.0	4.3	13.9%	3.2%	33.8%
11.5	4.5	19.2%	5.3%	39.7%
12.0	4.7	25.7%	8.4%	45.5%
12.5	4.9	33.2%	12.7%	69.6%
13.0	5.1	41.4%	18.3%	79.5%
13.5	5.3	50.0%	25.2%	91.0%
14.0	5.5	58.6%	33.4%	92.5%
14.5	5.7	66.8%	42.4%	94.0%
15.0	5.9	74.3%	51.9%	97.9%
15.5	6.1	80.8%	61.2%	98.4%
16.0	6.3	86.1%	70.0%	98.7%
16.5	6.5	90.4%	77.7%	98.9%
17.0	6.7	93.6%	84.1%	99.0%
17.5	6.9	95.9%	89.2%	99.2%
18.0	7.1	97.5%	93.0%	99.3%
18.5	7.3	98.5%	95.7%	99.5%
19.0	7.5	99.2%	97.5%	99.7%
19.5	7.7	99.5%	98.6%	99.8%
20.0	7.9	99.8%	99.2%	100.0%

Nota:

En estos casos la longitud del pene erecto fue medida desde la base del pene en el abdomen hasta la punta.

Fuentes:

1. *T. Schneider, H. Sperling, G. Lummen, et. al. Does penile size in younger men cause problems in condom use? a prospective measurement of penile dimensions in 111 young and 32 older men. Urology 2001 pp. 314-318.*

In 2001 Schneider et al, tested Korean men whose age ranged from 18-19 and 40-68 and measured an erect length of 14.48 cm for the former and 14.18 cm for the latter (14.48±5.7 cm, 14.18±5.58 cm).

2. *ANSELL RESEARCH - The Penis Size Survey (March 2001)*

A study conducted by LifeStyles Condoms found an average of 14.9 cm (5.9 in) with a standard deviation of 2.1 cm (0.8 in) (measured by staff). The purpose of this study was to ensure properly sized condoms were available. 401 college students volunteered to be measured during 2001 Spring Break in Cancún, Mexico, of which 300 gained an erection to be clinically measured (without pharmacological aid). The 300 effective measurements makes this study double the size of any prior study that uses medical staff to measure penis size.

Erect circumference.- A study conducted by LifeStyles Condoms during 2001 Spring Break in Cancún found an average of 12.6 cm (5.0 in) with a standard deviation of 1.3 cm (0.5 in).

Aumento Normal del Tamaño del Pene con la Edad

Edad (años)	10% lo tienen más chico que ...		50% lo tienen más chico que ...		90% lo tienen más chico que ...	
1	2.8 cm	1.1 in	4.2 cm	1.7 in	5.7 cm	2.2 in
2	3.3 cm	1.3 in	4.9 cm	1.9 in	6.2 cm	2.5 in
3	3.7 cm	1.5 in	5.3 cm	2.1 in	6.7 cm	2.6 in
4	4.0 cm	1.6 in	5.6 cm	2.2 in	7.0 cm	2.8 in
5	4.3 cm	1.7 in	5.8 cm	2.3 in	7.2 cm	2.8 in
6	4.6 cm	1.8 in	6.0 cm	2.3 in	7.3 cm	2.9 in
7	4.7 cm	1.8 in	6.1 cm	2.4 in	7.4 cm	2.9 in
8	4.8 cm	1.9 in	6.1 cm	2.4 in	7.4 cm	2.9 in
9	4.8 cm	1.9 in	6.1 cm	2.4 in	7.5 cm	2.9 in
10	4.8 cm	1.9 in	6.1 cm	2.4 in	7.8 cm	3.1 in
11	4.8 cm	1.9 in	6.2 cm	2.5 in	8.4 cm	3.3 in
12	5.1 cm	2.0 in	6.7 cm	2.6 in	9.4 cm	3.7 in
13	5.9 cm	2.3 in	7.9 cm	3.1 in	11.3 cm	4.5 in
14	7.3 cm	2.9 in	9.6 cm	3.8 in	13.2 cm	5.2 in
15	8.6 cm	3.4 in	10.8 cm	4.2 in	14.3 cm	5.6 in
16	9.6 cm	3.8 in	11.8 cm	4.6 in	14.8 cm	5.8 in
17	10.4 cm	4.1 in	12.6 cm	4.9 in	15.1 cm	5.9 in
18	10.6 cm	4.2 in	12.8 cm	5.0 in	15.2 cm	6.0 in

Nota:

Medir la longitud del pene totalmente erecto para obtener una medida precisa es difícil en un ambiente clínico. Algunos estudios preparan a los sujetos utilizando fármacos para inducir una erección, para lograr obtener resultados consistentes. Algunos otros estudios hacen las mediciones estirando el pene flácido tanto como sea cómodo. Esta tabla contiene datos utilizando este último método.

Fuentes:

Stretched Penile Length, measured from the pubo-penile skin junction to the tip of the glans (Shonfeld & Beebe. 1942, Journal of Urology, 48, 759-777).